arXiv:1208.1512v1 [cond-mat.stat-mech] 7 Aug 2012

General method for finding families of exact ground states of broad classes of classical Heisenberg models

Zhaoxi Xiong[1,*] and Xiao-Gang Wen[1,2]
[1]*Department of Physics, Massachusetts Institute of Technology, Cambridge, Massachusetts 02139, USA*
[2]*Perimeter Institute for Theoretical Physics, Waterloo, Ontario, N2L 2Y5 Canada*

(Dated: August 9, 2012)

We present a general method for systematically finding the families of exact ground states of classical Heisenberg models with the following properties: (i) they have the translational symmetries of infinite crystals; and (ii) they contain only bilinear terms, i.e. $\vec{S}_i \cdot \vec{S}_j$. With very limited exceptions, the method is applicable to all such models with one spin per unit cell, and models that have many spins per unit cell and satisfy certain conditions. Crystal dimensionality, range of coupling, etc. do not affect the validity. We also prove theorems that guarantee the existence of, in certain circumstances, a spiral state within the ground state manifold. Our formalism is predictive for disorder, and it might be generalized to higher-order terms (i.e. $(\vec{S}_i \cdot \vec{S}_j)^r$), anisotropic couplings, and arbitrary dimensional spins.

PACS numbers: 75.10.-b, 75.10.Hk

Exchange interactions dominate magnetic orders in solids. Despite the quantum nature, many studies on magnetic properties were set out classically. The reason, besides simplicity, is that classical models are indeed good approximations to quantum ones in certain cases, two of which are of large spins and of spin$-\frac{1}{2}$'s. The former has a footage in large-spin expansion [1], whereas the latter relies on the Hartree-level equivalence between spin$-\frac{1}{2}$'s and classical spins. Large localized spins appear quite commonly in rare earth materials. To achieve quantum anomalous Hall effect (QAHE) on them, a non-coplanar classical spin order is required so that the $\mathcal{T}$-reversal symmetry is broken [2]. On the other hand, in quest for quantum spin liquids (SL), investigations are often carried out as a quantum fluctuation analysis that succeeds classical calculations (see e.g. [3]). In these and many such ongoing fields, classical calculations play a fundamental and crucial role.

It is notoriously nontrivial to deduce the magnetic orders from a formally simple exchange interaction Hamiltonian. While numerous profound techniques have been developed in the quantum scenario (e.g. mean field theory [4], large-spin expansion [1], spin wave theory [5], high temperature expansion [6], transfer matrix methods [7, 8], and renormalization group theory [9]), less attention has been paid to classical models. To name a few, one prevalent idea is local minimization: instances are termwise [10] or cluster [11] minimization, as well as rewriting the energy in special local forms (see e.g. [12]). Another popular device is the classical spin wave theory (see e.g. [12, 13]), which usually assumes a spiral or helical ansatz (see Fig. 1) within which the energy is minimized. Furthermore, the Luttinger-Tisza (LT) methods first weaken the normalization constraints on spins and subsequently check the strong constraints (see e.g. [14]). There are also methods founded on some classical-quantum correspondences (see e.g. [15]). These, however, are not enough to give us a comprehensive understanding of classical spin orders, and a more satisfactory method is demanded.

In this Letter, we present a general method for finding the family of exact ground states of a classical exchange interaction Hamiltonian with the following properties: (i) the Hamiltonian has the translational symmetry of an infinite crystal; and (ii) the Hamiltonian contains only Heisenberg terms, i.e. $\vec{S}_i \cdot \vec{S}_j$ with the spins being real unit 3-vectors (these will be referred to as the two "basic assumptions"). We emphasize on the generality of the Hamiltonian: We did not make any assumption on the crystal dimensionality, the particular Bravais lattice, the internal structure of unit cells, or the range of exchange interactions. Thus it can be square, Kagome, face center cubic, or diamond structures with interactions among nearest-neighbors, second nearest-neighbors, and so on. We shall prove that, if the system has only one site (spin) in each unit cell, or if a system has many spins per cell but satisfies certain explicit conditions, then among the ground states there must exist a spiral state (see Fig. 1). We shall also show that, with fairly limited exceptions, our method can find *all* the ground states of *any* such system. Our "ground states" are not computed within any ansatz; namely, they are *rigorous*.

This Letter is organized as follows. We first focus on systems with one spin per unit cell; we shall prove the spiral theorem and give the guidelines for finding the whole ground state manifold (GSM). Then, with a little addi-



FIG. 1: Schematic (a) spiral and (b) helical states.

tional complication, we present a similar set of results for systems with multi spins per cell. Finally, we comment on the cases we cannot easily handle, and discuss a few other possible generalizations of the framework. It may be helpful to bear in mind that our method consists of only three essential steps: changing variables, figuring out the new restrictions, and determining the ground states.

One spin per unit cell.— A general classical exchange Hamiltonian in this case takes the form

$$H_{cl} = \frac{1}{2}\sum_{\vec{n}}\sum_{j} J_j \vec{S}_{\vec{n}} \cdot \vec{S}_{\vec{n}+\vec{\eta}_j}, \tag{1}$$

where real unit 3-vector $\vec{S}_{\vec{n}}$ is the classical spin at lattice point $\vec{n}$; $\vec{\eta}_j$ connects the two lattice points involved in an exchange term with parameter J_j; and the $\frac{1}{2}$ is due to double count. In this Letter we will keep the convention of *positive* signs before coupling constants. This problem is completely classical, but for convenience, we reformulate the energy expression with the help of a Hilbert space $\mathcal{H}$ that has an orthonormal basis $\{|\vec{n}\rangle\}$. Suppose there are N spins. Viewing $\vec{S}_{\vec{n}}$ as column vectors, we define the "ket state" and "bra state" corresponding to a particular spin configuration as

$$|\psi\rangle \equiv \frac{1}{\sqrt{N}}\sum_{\vec{n}} \vec{S}_{\vec{n}} \otimes |\vec{n}\rangle, \quad \langle\psi| \equiv \frac{1}{\sqrt{N}}\sum_{\vec{n}} \vec{S}^{\dagger}_{\vec{n}} \otimes \langle\vec{n}|, \tag{2}$$

with † representing the Hermitian conjugate of a matrix. We also define the effective Hilbert space Hamiltonian corresponding to the classical H_{cl} as

$$H \equiv \frac{1}{2}\sum_{\vec{n}}\sum_{j} J_j \mathbb{I} \otimes |\vec{n}\rangle\langle\vec{n}+\vec{\eta}_j|. \tag{3}$$

Then one figures that the quantity

$$\epsilon \equiv \langle\psi|H|\psi\rangle \tag{4}$$

recovers correctly the classical energy per spin. The bonus for defining H is that in $\mathcal{H}$, as an operator, H has an unambiguously defined set of eigenstates $|\vec{k}\rangle = \frac{1}{\sqrt{N}}\sum_{\vec{n}} e^{i\vec{k}\cdot\vec{n}}|\vec{n}\rangle$ and eigenvalues $\epsilon_{\vec{k}}$; we refer to the latter as the spectrum of H. One can also show from the basic assumptions the property $\epsilon_{-\vec{k}} = \epsilon_{\vec{k}}$. After change of basis, one finds Eq. (4) can be written in terms of $\epsilon_{\vec{k}}$ as

$$\epsilon = \sum_{\vec{k}} \epsilon_{\vec{k}} (\frac{1}{N}\vec{S}^{\dagger}_{\vec{k}}\vec{S}_{\vec{k}}), \tag{5}$$

with the new variables $\vec{S}_{\vec{k}}$ being the Fourier conjugates:

$$\vec{S}_{\vec{k}} \equiv \frac{1}{\sqrt{N}}\sum_{\vec{n}} \vec{S}_{\vec{n}} e^{-i\vec{k}\cdot\vec{n}}. \tag{6}$$

The above procedure formally looks as if we were defining quasi-particle excitations by Eq. (6) so as to diagonalize the Hamiltonian, but $\vec{S}_{\vec{k}}$ or $\vec{S}_{\vec{n}}$ are by no means annihilation operators. Now that we have a "diagonalized" form Eq. (5), we need to know what values $\{\vec{S}_{\vec{k}}\}$ can take. Noticing that the set of allowed $\{\vec{S}_{\vec{n}}\}$ is

$$\mathcal{D} = \big\{\{\vec{S}_{\vec{n}}\} | \vec{S}_{\vec{n}} \in \mathbb{R}^3,\ \vec{S}^{\dagger}_{\vec{n}}\vec{S}_{\vec{n}} = 1\big\}, \tag{7}$$

one can prove that a choice of $\{\vec{S}_{\vec{k}}\}$ is allowed *if and only if* it is contained in

$$\mathcal{I} = \big\{\{\vec{S}_{\vec{k}}\} | \vec{S}_{\vec{k}} \in \mathbb{C}^3, \vec{S}_{-\vec{k}} = \vec{S}^{*}_{\vec{k}}, \sum_{\vec{k}} \vec{S}^{\dagger}_{\vec{k}}\vec{S}_{\vec{k}+\vec{p}} = N\delta_{\vec{0}\vec{p}}, \forall\vec{p}\big\}. \tag{8}$$

$\mathcal{D}$ and $\mathcal{I}$ are the domain and the image of the map Eq. (6), respectively. The $\vec{p}$ in Eq. (8) is assumed to be located at, and the summations of $\vec{k}$ in Eqs. (5) (8) are over, the reciprocal lattice points in the Brillouin zone (BZ). In addition, in Eq. (8), $\vec{S}_{\vec{k}}$ is assumed to be periodic. Now we are ready to prove the following theorem.

Theorem 1 (Spiral ground state theorem (one-spin cells))**.** For any classical Heisenberg model that satisfies our two basic assumptions, and that has only one spin per unit cell, among the ground states there must exist a spiral state.

Proof. From the definition of $\mathcal{I}$ we know $\sum_{\vec{k}} \frac{1}{N}\vec{S}^{\dagger}_{\vec{k}}\vec{S}_{\vec{k}} = 1$; we also know $\frac{1}{N}\vec{S}^{\dagger}_{\vec{k}}\vec{S}_{\vec{k}} \geq 0$. Thus if $\vec{S}_{\vec{k}} \neq 0$ only at some global minima in the spectrum, then the resulting state must be a rigorous ground state. Suppose $\vec{k}_0$ is a global minimum; since $\epsilon_{-\vec{k}} = \epsilon_{\vec{k}}$, so is $-\vec{k}_0$. One can verify that the specification $\vec{S}_{\pm\vec{k}_0} = \frac{\sqrt{N}}{2}(1, \pm i, 0)^T$ for inequivalent $\pm\vec{k}_0$ (i.e., not connected by reciprocal translational vectors), or $\sqrt{N}(1,0,0)^T$ for equivalent $\pm\vec{k}_0$, is contained in $\mathcal{I}$ and is hence allowed. More explicitly, $\vec{S}_{\vec{n}} = (\cos(\vec{k}_0\cdot\vec{n}), -\sin(\vec{k}_0\cdot\vec{n}), 0)^T$ or $(e^{i\vec{k}_0\cdot\vec{n}}, 0, 0)^T$, for the two cases respectively. These are spiral states. □

From the proof one concludes that a sufficient and necessary condition for an allowed state to be a ground state is that its $\vec{S}_{\vec{k}}$ vanish at all not-the-lowest points, so as to have the same energy as the spiral state given by Theorem 1. This is the principle for finding the entire GSM: For any given Hamiltonian, we compute and examine the spectrum, write down equations according to $\mathcal{I}$ for $\vec{S}_{\vec{k}}$ at the minima, and determine the GSM. Moreover, since in this one-spin case, the distribution of spectral minima *completely* determines the GSM, it can be used for classification of all models. Be aware, however, that the distribution may be so special that the equations imposed by $\mathcal{I}$ are modified due to the periodicity of $\vec{S}_{\vec{k}}$ (e.g. Figs. 2(e)-(h)) or $\pm\vec{k}_i$ being equivalent (e.g. Figs. 2(a)-(d)).

We shall now classify two-dimensional (2D) systems in the afore-mentioned way and present their GSMs [16]. We are free (for any lattice) to assign integer numbers to n_x, n_y so that the BZ is $(-\pi,\pi]\times(-\pi,\pi]$ (we will use this



FIG. 2: For 2D systems with one spin per cell, the special distributions of spectral global minima, represented by the dots, in the BZ $(-\pi,\pi]\times(-\pi,\pi]$ (see text). The larger and the smaller dots are respectively defined as $\pm\vec{k}_0$ and $\pm\vec{k}_1$. This list is complete for the case of one pair of minima ((a) - (h)) but partial for the case of two pairs ((i) - (r)); definitely there can be more. In (o) - (r), $\pm k_1$ are generically located.

convention throughout the Letter). Special locations of the minima are partially enumerated in Fig. 2; the GSM of them and of some general cases are summarized in Table I. The expressions in Table I literally correspond to spiral phase, ferromagnetic (F)/antiferromagnetic (AF) phases, frustrated phases, spiral phase, frustrated phases, and helical phases (possibly with alternating helicities). Although the cases enumerated are obviously incomplete, one realizes that the GSM of any given Hamiltonian can be similarly worked out extremely easily, unless there are infinite number of spectral minima. Our formalism also makes phase transitions manifest, as there is usually an abrupt change in the number or locations of the lowest points when a transition occurs.

Let us apply our method to the J_1-J_2 models on square and triangular lattices, where $J_1,\ J_2$ correspond to nearest- and next nearest-neighbors respectively (recall our positive sign convention). The spectrum for square lattice is $\epsilon_{\vec{k}}=J_1(\cos k_x+\cos k_y)+2J_2\cos k_x\cos k_y$. The regimes with $J_1<0,\ J_2<0.5|J_1|$; $J_1>0,\ J_2<0.5J_1$; and $J_2>0.5$ respectively fall under Figs. 2(a)(d)(l), and hence have F, AF, and frustrated two-sublattice AF orders. The spectrum for triangular lattice (for some particular translational axes) is $\epsilon_{\vec{k}}=J_1[\cos k_x+\cos k_y+\cos(k_x-k_y)]+J_2[\cos(k_x+k_y)+\cos(2k_x-k_y)+\cos(2k_y-k_x)]$. The regime $J_1<0,\ J_2<\frac{1}{3}|J_1|$ is given by Fig. 2(a) and is ferromagnetic; for $J_1>0,\ J_2<\frac{1}{8}J_1$, there is a single pair of minima $\pm(\frac{2}{3}\pi,-\frac{2}{3}\pi)$, implying an angle $\frac{2}{3}\pi$ between neighboring spins; the remained cases have three pairs of lowest points, which are not included in Table I but are solvable.

Multi spins per cell.— For a general system with N unit cells that each has m spins, the Hamiltonian is of the form

$$H_{cl}=\frac{1}{2}\sum_{\vec{n}}\sum_{j}J_j\vec{S}_{\vec{n}a_j}\cdot\vec{S}_{\vec{n}+\vec{\eta}_jb_j},\tag{9}$$

TABLE I: The GSMs corresponding to different distributions of the spectral global minima, for 2D systems with one spin per cell. $\gamma\in\mathbb{R},\ \rho\in SO_3$ are arbitrary and preassigned.

Spectral global minima	GSM
Generic $\pm\vec{k}_0$	$\vec{S}_{\vec{n}}=\rho\begin{pmatrix}\cos(\vec{k}_0\cdot\vec{n})\\-\sin(\vec{k}_0\cdot\vec{n})\\0\end{pmatrix}$
Figs. 2(a)-2(d)	$\vec{S}_{\vec{n}}=\rho\begin{pmatrix}e^{i\vec{k}_0\cdot\vec{n}}\\0\\0\end{pmatrix}$
Figs. 2(e)-2(h)	$\vec{S}_{\vec{n}}=\rho\begin{pmatrix}\sqrt{2}\cos(\vec{k}_0\cdot\vec{n}+\frac{\pi}{4})\cos\gamma\\\sqrt{2}\sin(\vec{k}_0\cdot\vec{n}+\frac{\pi}{4})\sin\gamma\\0\end{pmatrix}$
Generic $\pm\vec{k}_0,\ \pm\vec{k}_1$	$\vec{S}_{\vec{n}}=\rho\begin{pmatrix}\cos(\vec{k}_i\cdot\vec{n})\\-\sin(\vec{k}_i\cdot\vec{n})\\0\end{pmatrix},\ i=0\text{ or }1$
Figs. 2(i)-2(n)	$\vec{S}_{\vec{n}}=\rho\begin{pmatrix}e^{i\vec{k}_0\cdot\vec{n}}\cos\gamma\\e^{i\vec{k}_1\cdot\vec{n}}\sin\gamma\\0\end{pmatrix}$
Figs. 2(o)-2(r)	$\vec{S}_{\vec{n}}=\rho\begin{pmatrix}\cos(\vec{k}_1\cdot\vec{n})\sin\gamma\\\sin(\vec{k}_1\cdot\vec{n})\sin\gamma\\e^{i\vec{k}_0\cdot\vec{n}}\cos\gamma\end{pmatrix}$

where $a_j,\ b_j=1,2,\ldots,m$ label the particular internal sites involved in a coupling term. Using an orthonormal basis $\{|\vec{n}a\rangle\}$ of a Hilbert space $\mathcal{H}$, we define

$$H\equiv\frac{1}{2}\sum_{\vec{n}}\sum_{j}J_j\mathbb{I}\otimes|\vec{n}a_j\rangle\langle\vec{n}+\vec{\eta}_jb_j|.\tag{10}$$

In the basis $|\vec{k}a\rangle\equiv\frac{1}{\sqrt{N}}\sum_{\vec{n}}e^{i\vec{k}\cdot\vec{n}}|\vec{n}a\rangle$, Eq. (10) can be written as $H=\sum_{\vec{k},a,b}\mathbb{I}\otimes|\vec{k}a\rangle\big(h(\vec{k})\big)_{ab}\langle\vec{k}b|$, where matrices $h(\vec{k})$ have well-defined orthonormal eigenvectors $\xi_{\vec{k}\alpha}$ and eigenvalues $\epsilon_{\vec{k}\alpha}$. In this multi-spin scenario, one observes a band-like structure, with α being the band label. Some universal properties can be proved: $h(-\vec{k})=h(\vec{k})^*$, $\epsilon_{-\vec{k}\alpha}=\epsilon_{\vec{k}\alpha}$, and $\xi_{-\vec{k},\alpha}=\xi^*_{\vec{k},\alpha}$ (convention). Following a similar procedure we find the per *unit cell* energy:

$$\epsilon=\sum_{\vec{k},\alpha}\epsilon_{\vec{k}\alpha}(\frac{1}{N}\vec{S}^\dagger_{\vec{k}\alpha}\vec{S}_{\vec{k}\alpha}),\tag{11}$$

with the new variables, in an abbreviate form, defined as

$$\mathscr{S}_{\vec{k}}\equiv(\frac{1}{\sqrt{N}}\sum_{\vec{n}}e^{-i\vec{k}\cdot\vec{n}}\mathscr{S}_{\vec{n}})\Xi_{\vec{k}},\tag{12}$$

where $\mathscr{S}_{\vec{n}}\equiv\big(\vec{S}_{\vec{n}1},\ldots,\vec{S}_{\vec{n}m}\big)$, $\mathscr{S}_{\vec{k}}\equiv\big(\vec{S}_{\vec{k}1},\ldots,\vec{S}_{\vec{k}m}\big)$, and $\Xi_{\vec{k}}\equiv\big(\xi_{\vec{k}1},\ldots,\xi_{\vec{k}m}\big)$. The image of the map Eq. (12) is also straightforwardly computable:

$$\begin{aligned}\mathcal{I}\ =\ &\{\{\vec{S}_{\vec{k}\alpha}\}|\vec{S}_{\vec{k}\alpha}\in\mathbb{C}^3,\ \vec{S}_{-\vec{k}\alpha}=\vec{S}^*_{\vec{k}\alpha},\\&\sum_{\vec{k}}(\Xi_{\vec{k}}\mathscr{S}^\dagger_{\vec{k}}\mathscr{S}_{\vec{k}+\vec{p}}\Xi^\dagger_{\vec{k}+\vec{p}})_{aa}=N\delta_{\vec{0}\vec{p}},\ \forall a,\vec{p}\},\end{aligned}\tag{13}$$

where the a on the second line are matrix indices. Now, we shall present a similar spiral theorem for multi-site unit cells but with an extra assumption, as below.

Theorem 2 (Spiral ground state theorem (multi-spin cells))**.** For any classical Heisenberg model with m spins per unit cell that satisfies the two basic assumptions, if there exist points $\pm\vec{k}_0$ and a band α_0 where $\epsilon_{\vec{k}\alpha}$ assumes the global minimum and such that $\xi_{\pm\vec{k}_0\alpha_0}$ can be chosen to take the form $\xi_{\pm\vec{k}_0\alpha_0} = \frac{1}{\sqrt{m}}\left(e^{\pm it_1}, \ldots, e^{\pm it_m}\right)^T$, then among the ground states there exists a spiral state.

The proof is analogous. The spiral state to be found is $\vec{S}_{\vec{n}a} = \left(\cos(\vec{k}_0 \cdot \vec{n} - t_a), -\sin(\vec{k}_0 \cdot \vec{n} - t_a), 0\right)^T$ or $\left(e^{i(\vec{k}_0 \cdot \vec{n} - t_a)}, 0, 0\right)^T$ for inequivalent or equivalent $\pm\vec{k}_0$ respectively. We further present two corollaries.

Corollary 1. For a system with two sites per unit cell, if the 2×2 matrices $h(\vec{k})$ contain no Pauli σ_z component for all $\vec{k}$, then there exists a spiral ground state.

Corollary 2. For a system with m spins per unit cell, if for every $\vec{k}$, $\left(h(\vec{k})\right)_{ab}$ only depends on $(a-b)$ mod m, then there exists a spiral ground state.

A classification in the multi-spin case should take into account the $\xi_{\vec{k}\alpha}$. Whenever a spiral ground state is guaranteed, the GSM can be determined by focusing only on the spectral minima. Unless there are infinite number of them, the calculation will be rather simple. It is fortunate that some models do fulfill the prerequisite on $\xi_{\vec{k}\alpha}$. For instance, Corollary 1 is applicable to the honeycomb $J_1 - J_2 - J_3$ model, which has been studied by classical spin waves [12, 13]; our method can determine the ground state energy and, in regimes with finite number of minima, the whole GSM. Another instance is the checkerboard $J_1 - J_2$ model, where J_1, J_2 correspond to straight and diagonal edges respectively (see e.g. [17]), and whose $h(\vec{k})$ has no σ_z component along $|\vec{k}_x| = |\vec{k}_y|$. For $\frac{J_2}{|J_1|} < 1$, a single minimum appears at $\vec{k} = 0$, implying a F ($J_1 < 0$) or AF ($J_1 > 0$) phase. For $\frac{J_2}{|J_1|} > 1$, the minima cover the entire BZ boundary.

Discussion.— We cannot easily handle the following cases: the case of an infinite number of spectral minima, and the multi-spin case in which the $\xi_{\vec{k}\alpha}$ condition in Theorem 2 is not satisfied. Nevertheless, one can still gain some knowledge within our framework. For example, ideally QAHE desires a purely classical Hamiltonian with *no* coplanar ground state, and our results imply such a system must go beyond the spiral theorems. Furthermore, since the dimension of GSM is not greater than the number of independent variables (of the same order as the number of nonzero $\vec{S}_{\vec{k}\alpha}$), a disordered system with an infinite dimensional GSM must have infinite number of spectral minima or violate the $\xi_{\vec{k}\alpha}$ condition.

There exist a few possible generalizations of the framework. To accommodate Hamiltonians with exclusive higher-order terms, i.e. $(\vec{S}_i \cdot \vec{S}_j)^r$, one may redefine $|\psi\rangle \equiv \frac{1}{\sqrt{N}} \sum_{\vec{n},a} (\vec{S}_{\vec{n}a})^{\otimes r} \otimes |\vec{n}a\rangle$. To include anisotropic couplings of the form $\vec{S}_i^{\mu} J^{\mu\nu} \vec{S}_j^{\nu}$, one may replace the $J_j \mathbb{I}$ in Eq. (10) with a matrix J_j. In addition, the spins can be generalized to arbitrary dimension, adapting the formalism to XY and Ising models, etc.

Upon completion of this work, we noticed the similar conclusions about spiral states given by the LT methods (see e.g. [14]), but we have derived them in a different and more general framework. In this framework, it is rather simple to determine the entire GSM, classify all models, predict phase transitions, and draw hints for noncoplanarity and disorder. Our philosophy of keeping the full information, as opposed to the LT methods, seems so natural that it should have been discovered decades ago.

Z. Xiong acknowledges Q. Lin for helpful comments. This work is supported by NSF Grant No. DMR-1005541 and NSFC 11074140.

---

[*] Electronic address: xiong@mit.edu

[1] T. Holstein and H. Primakoff, Phys. Rev. **58**, 1098 (1940).

[2] J. Ye *et al.*, Phys. Rev. Lett. **83**, 3737 (1999); K. Ohgushi, S. Murakami, and N. Nagaosa, Phys. Rev. B **62**, R6065 (2000).

[3] P. Chandra and B. Doucot, Phys. Rev. B **38**, 9335 (1988).

[4] See for example, G. Grosso and G. P. Parravicini, *Solid State Physics* (Academic, San Diego, 2003), pp. 620-627.

[5] F. Bloch, Z. Phys. **61**, 206 (1930); P. W. Anderson, Phys. Rev. **86**, 694 (1952); R. Kubo, *ibid.* **87**, 568 (1952).

[6] See for example, *Phase Transitions and Critical Phenomena*, edited by C. Domb and M. S. Green (Academic, London, 1974), Vol. 3.

[7] E. Ising, Z. Phys. **31**, 253 (1925).

[8] L. Onsager, Phys. Rev. **65**, 117 (1944).

[9] K. G. Wilson, Rev. Mod. Phys. **47**, 773 (1975).

[10] C. A. M. Mulder, H. W. Capel, and J. H. H. Perk, Physica (Utrecht) **112B**, 147 (1982).

[11] D. H. Lyons and T. A. Kaplan, J. Phys. Chem. Solids **25**, 645 (1964).

[12] J. B. Fouet, P. Sindzingre, and C. Lhuillier, Eur. Phys. J. B **20**, 241 (2001).

[13] E. Rastelli, A. Tassi, and L. Reatto, Physica (Utrecht) **97B**, 1 (1979).

[14] T. A. Kaplan and N. Menyuk, Philos. Mag. **87**, 3711 (2007).

[15] M. Van den Nest, W. Dür, and H. J. Briegel, Phys. Rev. Lett. **98**, 117207 (2007).

[16] See Supplemental Material at [to be specified] for exemplifying calculations.

[17] B. Canals, Phys. Rev. B **65**, 184408 (2002).

arXiv:1208.1512v1 [cond-mat.stat-mech] 7 Aug 2012

Supplemental Material for "General method for finding families of exact ground states of broad classes of classical Heisenberg models"

Zhaoxi Xiong[1,*] and Xiao-Gang Wen[1,2]
[1] Department of Physics, Massachusetts Institute of Technology, Cambridge, Massachusetts 02139, USA
[2] Perimeter Institute for Theoretical Physics, Waterloo, Ontario, N2L 2Y5 Canada

(Dated: August 9, 2012)

In this Supplemental Material we show explicitly how to determine the entire ground state manifold (GSM) of a system within the framework we have developed. We consider two exemplifying cases for two-dimensional systems with one spin per cell: (i) the case in which there is a single pair of generically located minima $\pm\vec{k}_0$ in the spectrum; and (ii) the case in which there are two specially located minima $\vec{k}_0 = (0,0)$, $\vec{k}_1 = (\pi, 0)$ in the spectrum.

Consider case (i), which corresponds to the first item in Table I of the main paper. In this case, any ground state can have nonzero $\vec{S}_{\vec{k}}$ only at $\pm\vec{k}_0$. Hence, we set

$$\vec{S}_{\vec{k}_0} = \sqrt{\frac{N}{2}} \begin{pmatrix} x \\ y \\ z \end{pmatrix}, \quad \vec{S}_{-\vec{k}_0} = \sqrt{\frac{N}{2}} \begin{pmatrix} x^* \\ y^* \\ z^* \end{pmatrix}, \tag{1}$$

and zero otherwise. We must check if such a specification is contained in

$$\mathcal{I} = \big\{\{\vec{S}_{\vec{k}}\} | \vec{S}_{\vec{k}} \in \mathbb{C}^3,\ \vec{S}_{-\vec{k}} = \vec{S}^*_{\vec{k}},\ \sum_{\vec{k}} \vec{S}^\dagger_{\vec{k}} \vec{S}_{\vec{k}+\vec{p}} = N\delta_{\vec{0}\vec{p}}, \forall \vec{p}\big\}. \tag{2}$$

The first two conditions are fulfilled. The third is effective when $\vec{p} = 0$ and $\pm 2\vec{k}_0$ (up to a reciprocal translational vector). The cases of $\vec{p} = 0,\ 2\vec{k}_0$ respectively require

$$|x|^2 + |y|^2 + |z|^2 = 1, \quad x^2 + y^2 + z^2 = 0. \tag{3}$$

The case of $\vec{p} = -2\vec{k}_0$ gives a condition conjugate to that of $\vec{p} = 2\vec{k}_0$. The set of solutions to Eq. (3) would give the GSM. Written in terms of real and imaginary parts, namely $x \equiv x' + ix''$, $y \equiv y' + iy''$, $z \equiv z' + iz''$, Eq. (3) reduces to

$$||(x', y', z')|| = ||(x'', y'', z'')|| = \frac{1}{\sqrt{2}}, \quad (x', y', z') \perp (x'', y'', z''), \tag{4}$$

where we have viewed (x', y', z'), (x'', y'', z'') as 3-component vectors. Now, apparently the set of solutions to Eq. (4) is, up to an arbitrary rotation,

$$(x', y', z') = (\frac{1}{\sqrt{2}}, 0, 0), \quad (x'', y'', z'') = (0, \frac{1}{\sqrt{2}}, 0), \tag{5}$$

whence

$$\vec{S}_{\pm\vec{k}_0} = \frac{\sqrt{N}}{2} \begin{pmatrix} 1 \\ \pm i \\ 0 \end{pmatrix}. \tag{6}$$

The GSM is therefore

$$\vec{S}_{\vec{n}} = \rho \begin{pmatrix} \cos(\vec{k}_0 \cdot \vec{n}) \\ -\sin(\vec{k}_0 \cdot \vec{n}) \\ 0 \end{pmatrix}, \tag{7}$$

where $\rho \in SO_3$ is an arbitrary rotation.

---

[*] Electronic address: xiong@mit.edu

Next, consider case (ii), which falls under the fifth item in Table I of the main paper. The periodicity and the definition of $\mathcal{I}$ together imply that $\vec{S}_{\vec{k}_0},\ \vec{S}_{\vec{k}_1}$ are real. We thus set

$$\vec{S}_{\vec{k}_0} = \sqrt{\frac{N}{2}} \begin{pmatrix} x \\ y \\ z \end{pmatrix}, \quad \vec{S}_{\vec{k}_1} = \sqrt{\frac{N}{2}} \begin{pmatrix} u \\ v \\ w \end{pmatrix}, \tag{8}$$

where $x,\ y,\ z,\ u,\ v,\ w \in \mathbb{R}$. The $\vec{p} = (0,0),\ (\pi,0)$ cases of $\sum_{\vec{k}} \vec{S}^{\dagger}_{\vec{k}} \vec{S}_{\vec{k}+\vec{p}} = N\delta_{\vec{0}\vec{p}}$ give two independent constraints:

$$||(x,y,z)||^2 + ||(u,v,w)||^2 = 2, \quad (x,y,z)\cdot(u,v,w) = 0. \tag{9}$$

The solutions are, up to an arbitrary rotation,

$$(x,y,z) = \sqrt{2}(\cos\gamma, 0, 0), \quad (u,v,w) = \sqrt{2}(0, \sin\gamma, 0), \tag{10}$$

where $\gamma \in \mathbb{R}$ is another arbitrary parameter. Consequently,

$$\vec{S}_{\vec{k}_0} = \sqrt{N} \begin{pmatrix} \cos\gamma \\ 0 \\ 0 \end{pmatrix}, \quad \vec{S}_{\vec{k}_1} = \sqrt{N} \begin{pmatrix} 0 \\ \sin\gamma \\ 0 \end{pmatrix}. \tag{11}$$

Therefore, the GSM is

$$\vec{S}_{\vec{n}} = \rho \begin{pmatrix} \cos\gamma \\ (-1)^{n_x}\sin\gamma \\ 0 \end{pmatrix}, \tag{12}$$

where $\rho \in SO_3,\ \gamma \in \mathbb{R}$ are arbitrary.